




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 35/78</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/62532</b>
			(43) International Publication Date: 9 December 1999 (09.12.99)
(21) International Application Number: PCT/US99/11880 (22) International Filing Date: 28 May 1999 (28.05.99)  (30) Priority Data: 60/087,680                      2 June 1998 (02.06.98)                      US 09/311,751                      14 May 1999 (14.05.99)                      US  (71) Applicant: BAYLOR COLLEGE OF MEDICINE [US/US]; One Baylor Plaza, Houston, TX 77030 (US).  (72) Inventors: MARKAVERICH, Barry, M.; 112 West Rainbow Ridge Circle, The Woodlands, TX 77380 (US). FAITH, Robert; 9623 South Petersham, Houston, TX 77031 (US). MANI, Sheila; 3416 Rice Boulevard, Houston, TX 77005 (US).  (74) Agents: REMENICK, James et al.; Baker & Botts, L.L.P., The Wamer, Suite 1300, 1299 Pennsylvania Avenue, N.W., Washington, DC 20004-2400 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published <i>With international search report.</i>	

(54) Title: CORN EXTRACT CONTRACEPTIVE

## (57) Abstract

Novel compositions extracted from corn products provide contraceptive and anti-neoplastic activities. Using novel extraction procedures, compositions may be isolated from corn products. For example, *Zea mays* plant product is extracted in a first solvent to produce a solvent extract, the solvent extract is dried to produce an extracted solid, the extracted solid is solubilized in a second solvent, the solubilized extract is purified in a chromatographic process, and an active fraction is collected from the chromatographic process. Compositions may be applied to animal bedding or food, and are adaptable to any suitable method of administration. The contraceptive activity of these compositions is effective for both males and females.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CORN EXTRACT CONTRACEPTIVE

### Background of the Invention

#### 1. Field of the Invention

This invention relates to pharmaceutical compositions derived from corn  
5 extracts useful as contraceptive compositions and antineoplastic compositions.

#### 2. Description of the Background

Contraception is often desired to control reproduction. While a plurality  
of contraceptive compositions and methods are available commercially, most methods  
of chemical contraceptives are active through a limited number of pathways. Thus, the  
10 majority of contraceptive compositions still involve the use of one or more natural or  
synthetic hormones such as estrogen, progestin and ethinyl estradiol.

While natural and synthetic hormones are effective, their use has not  
been without substantial and undesirable side effects. Numerous side effects have been  
associated with the use of natural or synthetic hormones in contraceptives, such as, for  
15 example, thrombophlebitis, thrombosis, pulmonary embolism, coronary thrombosis,  
myocardial infarction, cerebral thrombosis, cerebral hemorrhage, hypertension, irregular  
bleeding, and irregular menstrual cycle.

Even with the reduction (but not elimination) of side effects through  
improved formulation, a number of disadvantages are still associated with current  
20 contraceptive compositions. Oral contraceptives rely on a high degree of patient  
compliance. For example, the risk of pregnancy increases with each pill missed or each  
pill taken out of order. Also, implantable contraceptives have many undesirable side  
effects and enjoy limited consumer acceptance. Another disadvantage of conventional  
contraceptives is that they are only effective when taken by females. Thus, the type of  
25 contraceptive choices available to couples is limited.

### Summary of the Invention

The present invention overcomes many of the limitations, problems and  
disadvantages associated with current strategies and designs for contraception and  
provides compositions and methods for contraception. As embodied and broadly  
30 described herein, the present invention is directed to a corn extract, such as an extract

- 2 -

from corn cob, useful as a contraceptive. In addition, extracts of the present invention are also useful as antineoplastic agents.

One embodiment of the invention is directed to a process for producing a therapeutic substance. In this process, a *Zea mays* plant product is extracted to produce a solvent extract, the solvent extract is dried to produce an extracted solid, the  
5 extracted solid is resolubilized in a solvent to form a resolubilized extract, the resolubilized extract is purified in a chromatography process, such as a reverse phase column chromatography process, and an active fraction is collected from the chromatography process. The *Zea mays* plant product used for starting material may  
10 be corn kernel, corn cob, animal bedding material made from corn cob, and mixtures of these materials.

Another embodiment of the invention is directed to a therapeutic substance produced by the process of the invention. The therapeutic substance may be in the form of a salt, a compound, a complex or a combination of these forms.

15 Another embodiment of the invention is directed to a pharmaceutical preparation that contains the therapeutic substance produced by the process of this invention. The pharmaceutical preparation may include, in addition to the active ingredients, a pharmaceutically acceptable carrier.

Another embodiment of the invention is directed to a composition for  
20 effecting contraception in a patient. The composition is preferably made using the process of the invention. The patient may be any mammal such as, for example, humans, monkeys, mice, rats, guinea pigs, rabbits, hamsters, horses, cattle, pigs, sheep, goats, dogs and cats.

Another embodiment of the invention is directed to a method of  
25 contraception comprising isolating an extract from a *Zea mays* plant product using a polar solvent and administering an effective amount of the extract to an individual to effect contraception.

Another embodiment of the invention is directed to a method for treating a patient with a neoplastic disorder comprising administering to the patient an  
30 antineoplastic composition comprising an effective quantity of the therapeutic

- 3 -

compound of the invention. The antineoplastic composition of the invention may further comprise an antineoplastic compound or agent, which may be, for example, methotrexate.

Another embodiment of the invention is directed to a method of  
5 inhibiting neoplastic activity comprising the steps of isolating an antineoplastic extract from a *Zea mays* plant product using a polar solvent and administering an effective amount of the antineoplastic extract to an individual. The method may further comprise the step of administering an antineoplastic compound, such as methotrexate.

Another embodiment of the invention is directed to an isolated and  
10 purified compound derived from an extract of *Zea mays* having a molecular weight of 348 amu. Preferably, the compound is isolated from extract that is derived from *Zea mays* corn cob using a polar solvent.

Another embodiment of the invention is directed to an isolated and  
purified compound derived from an extract of *Zea mays* having a molecular weight of  
15 353 amu. Preferably, the compound is isolated from extract that is derived from *Zea mays* corn cob using a polar solvent.

Another embodiment is directed to an isolated and purified compound  
derived from an extract of a *Zea mays* plant product, the extract being derived from the  
*Zea mays* plant product according to the methods of the invention, the compound  
20 eluting as a single UV-absorption peak from a C<sub>18</sub> reversed phase HPLC column having a flow rate of 2 ml per minute, as described below, the compound eluting approximately 56 minutes after injection of the extract sample onto the column, and having a molecular weight of either 348 amu or 353 amu.

Another embodiment is directed to an isolated and purified compound  
25 derived from an extract of a *Zea mays* plant product, the extract being derived from the *Zea mays* plant product according to the methods of the invention, the compound eluting as a single UV-absorption peak from a C<sub>18</sub> reversed phase HPLC column having a flow rate of 2 ml per minute, as described below, the compound eluting approximately 60 minutes after injection of the extract sample onto the column, and having a  
30 molecular weight of either 348 amu or 353 amu.

Other embodiments and advantages of the invention are set forth, in part, in the description which follows and, in part, will be obvious from this description and may be learned from the practice of the invention.

### Description of the Drawings

- |    |             |  |
|----|-------------|--|
| 5  | Figure 1    | Uterine wet weight and DNA content of ovariectomized rats housed on wire bedding or estrogenic substance-containing wood chip bedding for 14 days. |
|    | Figure 2(A) | Uterine wet weight of ovariectomized and non-ovariectomized rats housed on wire or GCCB bedding for 12 days.                                       |
| 10 | Figure 2(B) | Uterine DNA content of ovariectomized and non-ovariectomized rats housed on wire or GCCB bedding for 12 days.                                      |
|    | Figure 3    | Percentage of rats undergoing normal estrous cycles when housed on wire or corn cob bedding for two and five week periods.                         |
|    | Figure 4    | Percentage of rats undergoing normal estrous cycles when housed on wire, corn cob or transferred from corn cob to wire bedding.                    |
| 15 | Figure 5(A) | Effects of housing conditions on female sexual behavior.   |
|    | Figure 5(B) | Effects of housing conditions on male sexual behavior.   |
|    | Figure 6(A) | Effects of corn cob extracts on estrogen receptor binding by estradiol.  |
|    | Figure 6(B) | Effects of corn cob extracts on Type II site binding by estradiol.   |
| 20 | Figure 7    | Effects of corn cob extract and estradiol on cell growth.  |
|    | Figure 8    | Effects of corn cob extract column fractions on cell growth.   |
|    | Figure 9    | Effects of corn cob extract on estrogen receptor binding and Type II site binding by estradiol.  |
|    | Figure 10   | Effects of corn cob extract column fractions on cell growth.   |
| 25 | Figure 11   | Effects of ground corn cob bedding extract on estrogen receptor-mediated gene transcription.   |
|    | Figure 12   | Effects of antiestrogen on the stimulation of cell proliferation by estrogen or the purified corn cob bedding extract.                             |
|    | Figure 13   | Effects of various doses of corn cob extract on rat estrous cycles.  |
| 30 | Figure 14   | Effects of diluted corn cob extract on cells lacking estrogen receptors.   |

- 5 -

- Figure 15 Effects of various concentrations of diluted corn cob bedding extract on proliferation of cells lacking estrogen receptors.
- Figure 16 Bedding extract effects on Methotrexate inhibition of MCF-7 breast cancer cells.
- 5 Figure 17 Effect of extract from fresh corn on cell proliferation.
- Figure 18A Purification of mitogenic agent(s) from GCCB extract.
- Figure 18B Re-chromatography of mitogenic agent(s) from GCCB extract.
- Figure 19 Molecular weight determination of a putative mitogenic agent(s) in corn cob bedding extract.
- 10 Figure 20 Molecular weight determination of a putative mitogenic agent(s) in corn cob bedding extract.
- Figure 21 HPLC purification of a corn cob mitogen.

### Description of the Invention

This invention is directed to corn extracts and uses of compositions  
15 derived from corn extracts. Corn, also known as *Zea mays*, maize, or maiz, may be any of numerous cultivated forms of usually tall annual cereal grass bearing grains or kernels on large ears.

The present invention is also directed to a process for producing a  
therapeutic substance from a *Zea mays* plant product. The therapeutic substance may  
20 be a substance that shows contraceptive activity or antineoplastic activity or a combination of both activities. Compositions of the invention are useful for the control of sexual behavior in both males and females and may function as a chemical substitute for surgical castration. Potential uses of the invention include its use in a composition for blocking lordosis, an environmentally safe contraceptive for pest population control  
25 and a food additive with contraceptive activity.

Contraceptive activity may be any activity that diminishes the likelihood of, or prevents conception. Contraceptives may work through many mechanisms including: (1) inhibiting the estrous cycle in females; (2) inhibiting estrus (the recurrent, restricted period of sexual receptivity in female mammals, other than humans,  
30 characterized by intense sexual urge); (3) blocking ovulation or prevention of normal

- 6 -

egg formation in the female; (4) the reduction of sexual desire or behavior in the male or female; (5) blocking, reducing, or preventing formation of normal sperm in the male; (6) inducing erectile dysfunction in the male; and (7) inhibiting post fertilization implantation of the zygote. Further, contraceptives may work by a combination of any  
5 of the above mechanisms.

Antineoplastic activity may be any activity that inhibits or prevents the development of neoplasms or any activity that would check the maturation or proliferation of malignant cells. Inhibition of neoplasm may be due to an anti-hypertrophic (the prevention of the enlargement or overgrowth of an organ or part due  
10 to an increase in size of its constituent cells) or anti-hyperplastic (the prevention of abnormal multiplication or increase in the number of normal cells in normal arrangement in a tissue) activity as the antineoplastic activity.

The antineoplastic activity may be a direct result of the therapeutic substance of the *Zea mays* plant product, or alternately, the antineoplastic activity may  
15 be only present when the therapeutic substance of *Zea mays* is combined with another therapeutic substance. For example, a therapeutic substance may selectively induce the growth of neoplastic cells such that they are more susceptible to the effects of antimetogenic chemotherapeutic substances.

The starting material for extraction of the therapeutic substance is *Zea*  
20 *mays*, which is also known as poaceae, corn, maize, mealie and indian corn. *Zea mays* refers to any annual grass of the grass order *Cyperales*, family *Zea* and genus *mays* and includes all cultivars in the *mays* family such as *everta* (popcorn), *indurata* (flint corn), *indents* (dent corn), *amylacea* (flour corn), *saccharata* (soft corn, sweet corn) and *tunicata* (pod corn). Corn kernel or corn seed are the common names for what is  
25 technically the fruit of *Zae mays*. The *Zea mays* plant part or product which may be used for extraction may include, for example, the corn cob or the corn kernel. The starting material may be fresh, frozen, dried, or preserved *Zea mays* plant parts or products and mixtures thereof. Dried forms of *Zea mays* plant product include dried ground corn cob bedding (GCCB) which is commercially sold under the trade name



- 7 -

BEDOCOB™. The animal bedding material may be any commercially available ground corn cob bedding material.

The therapeutic substance may be extracted from the *Zea mays* plant parts or products by solvent extraction followed by drying to produce an extracted solid.

- 5 The extracted solid may be resolubilized in solvent before use. The solvent used for extraction and solubilization may be the same. Alternatively, the extracted solid may be further purified by reverse phase chromatography. In reverse phase chromatography, fractions eluted from a column are collected and tested for activity.

- 10 Solvent extraction may be performed using any primary, secondary and tertiary alcoholic solvent such as methanol, ethanol, propanol, butanol, sec butyl alcohol and mixtures of these alcohols. The solvent used for extraction may also be selected from the group consisting of alcohols, ethers, ketones, water and mixtures thereof. In a preferred embodiment, the solvent is the alcohol methanol.

- 15 Extraction may be performed using standard extraction procedures such as those listed in the Examples section or in Protein Purification (R.K. Scopes, 2nd edition, Springer-Verlag, New York, 1987). The extraction may be performed under room temperature or at a temperature of between about 0° and about 100°C and may be performed for about 1 hour to about 1 month. In one embodiment of the invention, corn cob, corn kernel or corn on the cob (or mixtures thereof) is immersed in an alcohol  
20 such as methanol in an enclosed chamber for a period of time of between one hour to one month. The starting material may be whole or may be processed to a small size more convenient for extraction. The temperature of the alcohol may be controlled between 0°C and 100°C.

- 25 After extraction, the liquid phase of the extraction is separated from the solid phase. Separation may be by any commonly known method such as filtering, settling, and centrifugation. After the separation, the mixture may be dried to form a dried extract. Drying may be performed by vacuum evaporation such as in a centrifugal vacuum evaporator, by exposure to dry air such as dried nitrogen, by lyophilization, or by evaporation with or without the assistance of heat. The solid extract may be

- 8 -

resolubilized in a suitable solvent such as, for example, dimethyl sulfoxide (DMSO), methanol, water or mixtures of these chemicals.

One preferred purification method for the purification of the therapeutic substance is reverse phase chromatography including reverse phase high performance liquid chromatography (reverse phase HPLC). Reverse-phase HPLC may be performed using hydrophobic adsorbents such as short-chain aliphatic groups attached to the matrix, and increasing organic solvent concentrations to elute. A preferred hydrophobic absorbent is a C<sub>18</sub> solid phase and a preferred hydrophilic solvent is a mixture of methanol and water. In a preferred embodiment, the method of reverse phase chromatography uses a C<sub>18</sub> solid phase column and the active fraction is eluted from the column with a solution of about 80% methanol and about 20% water.

Chromatography may be performed in column or batch mode. In column chromatography, a sample is loaded onto the column and eluted with successively higher concentrations of a hydrophilic solvent. The increase in solvent hydrophobicity may be continuous, such as, for example, from 20% to 100% methanol in 100 ml of chromatography liquid. Alternatively, the increase in solvent hydrophobicity may be discrete, such as, for example, loading with water and 20% methanol, elution with 40% methanol in water, 60% methanol in water, 80% methanol in water and 100% methanol. Other purification methods which may be used in addition to reverse phase chromatography include low pressure column chromatography, batch chromatography, precipitation, specific adsorbent chromatography, gel filtration, HPLC and combinations of these methods.

In a preferred embodiment of the invention, the reverse phase column is loaded with 20% methanol in water and washed with 60% methanol in water and a fraction showing the desired contraceptive and antineoplastic activity is eluted with 80% methanol in water. While one elution condition has been disclosed, other elution conditions may be determined by one of ordinary skill in the art without undue experimentation. For example, a C<sub>8</sub> to C<sub>18</sub> column, such as columns like a C<sub>9</sub>, C<sub>10</sub>, C<sub>11</sub>, C<sub>12</sub>, C<sub>13</sub>, C<sub>14</sub>, C<sub>15</sub>, C<sub>16</sub> or C<sub>17</sub> column may require different elution conditions than a C<sub>18</sub> column.

- 9 -

Another embodiment of the invention is directed to a therapeutic substance produced by the process of the invention. The therapeutic substance may have contraceptive or antineoplastic activities and may be in a form of a salt, a compound, a complex or a combination of these forms.

5 Another embodiment of the invention is directed to a pharmaceutical preparation comprising a therapeutic substance produced by the process of the invention with a pharmaceutically acceptable carrier. The pharmaceutical preparation may be in the form of an aqueous solution for parenteral and oral use, a capsule for oral use, or in other forms such as unguents, ointments, creams, suppositories and ovules.  
10 Administration routes may include parenteral, oral, ocular, periodontal, rectal or vaginal delivery, among others.

Another embodiment of the invention is directed to a composition for effecting contraception in a patient. The composition includes a contraceptive substance produced by the process of this invention. The patient may be a male or  
15 female mammal and may belong to any mammalian group including, but not limited to, hominids (humans), bovine, porcine, equine, caprine, canine, feline, murine and ovine groups. The methods and compositions of the present invention are useful for all types of laboratory animals, including those of the orders Rodentia (*e.g.* mice, rats, hamsters, guinea pigs, gerbils) Lagormorpha (*e.g.* rabbits) and Primates.

20 The contraceptive composition may be administered to a mammal by daily administration, weekly administration, monthly administration, yearly administration, continuous administration and cyclic administration. Methods of administration may be in the form of a pill or liquid capsule, an injection, a patch, a silastic implant or any other suitable means known in the art.

25 Another embodiment of the invention is directed to a method of contraception comprising isolating an extract from a *Zea mays* plant product using a polar solvent and administering an effective amount of the extract to an individual to effect contraception. The plant product is preferably selected from the group consisting of corn kernel, corn cob, animal bedding material made from corn cob, and mixtures  
30 thereof. The step of isolating the extract may further comprise the step of purifying the

- 10 -

extract by a chromatographic process before administration. In a preferred embodiment, the chromatographic process is reverse phase chromatography and the polar solvent is an alcohol.

The contraceptives and methods of contraception of the present invention may be used on both females and males. In females, contraception may be effected by inhibiting estrus, inhibiting the estrous cycle, inhibiting or blocking ovulation, inhibiting or blocking oogenesis, inhibiting or reducing sexual desire, inhibiting zygote implantation or combinations thereof. In males, contraception may be effected by inhibiting or reducing sexual desire, inhibiting, blocking, reducing or preventing spermatogenesis, inhibiting erectile function, or combinations thereof. The invention may be used on any male or female mammal selected from the group consisting of humans, monkeys, mice, rats, guinea pigs, rabbits, hamsters, horses, cattle, pigs, sheep, goats, dogs and cats. The extract may be administered to the individual by any suitable route of administration, including, but not limited to, orally, topically, intramuscularly, intravenously, subcutaneously, vaginally, rectally or by pulmonary absorption.

Another method of the invention is directed to a method for treating a patient with a neoplastic disorder comprising administering to the patient an antineoplastic composition which contains an effective quantity of the antineoplastic therapeutic compound of the present invention. The antineoplastic composition may further comprise an antineoplastic compound or agent, which may be, for example, methotrexate.

While all neoplasms may be treated with the therapeutic compounds of the invention, preferred neoplasms may be those that are responsive to antineoplastic agents, mitogen therapy or cell cycle specific agents such. These agents and mitogens may be used alone or in combination. Antineoplastic agents are known to those of skill in the art. Examples of antineoplastic agents which may be used alone or in combination include 5-fluorouracil, 6-mercaptopurine, 6-thioguanine, alkylating agents, androgens, antiadrenals, antiandrogens, antiestrogens, antimetabolites, asparaginase, bleomycin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, corticosteroids,

- 11 -

cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin, doxorubicin, estrogens, etoposide, fludarabine, fluorouracil, flutamide, hexamethylmelamine, hormones, hydroxyurea, hypothalamic agents, idarubicin, ifosfamide, intrapleural, lomustine, mechlorethamine, melphalan, methotrexate, mithramycin, mitomycin, mitotic inhibitors, mitoxantrone, paclitaxel, procarbazine, progestins, streptozocin, taxol, thiotepa, vinblastine, vincristine, vinorelbine and combinations and analogs thereof. Of these agents at least vincristine, vinblastine, vinorelbine, etoposide, paclitaxel, methotrexate, 6-mercaptopurine, 6-thioguanine, fludarabine, 5-fluorouracil and cytarabine may be considered cell cycle specific agents. In a preferred embodiment, the antineoplastic compound is methotrexate.

Neoplasms which may be treated may be any cancer that responds to mitogen therapy or cell cycle specific agents such as: acute lymphocytic leukemia; non Hodgkin's lymphoma; and breast, testicular, gastrointestinal tract, lung, skin, head, and neck tumors. Other neoplasms which may be treated include bone cancer, liver cancer and pancreatic cancer, prostate cancer, lymphomas and testicular cancer.

Another embodiment of the invention is directed to a method of inhibiting neoplastic activity comprising the steps of isolating an antineoplastic extract from a *Zea mays* plant product using a polar solvent and administering an effective amount of the antineoplastic extract to an individual. The method may further comprise the step of administering an antineoplastic compound such as methotrexate. In a preferred embodiment, the neoplastic activity to be inhibited is breast cancer cell proliferation. The extract may be administered by any of numerous methods such as, for example, orally, topically, intramuscularly, intravenously, subcutaneously, vaginally, rectally or by pulmonary absorption.

Another embodiment of the invention is directed to an isolated and purified compound derived from an extract of *Zea mays*. A *Zea mays* extract is prepared according to the methods of the present invention. The isolated and purified compound is then purified from the extract using a semipreparative C<sub>18</sub> reversed phase HPLC column equilibrated in 70% water and 30% acetonitrile containing 0.05% acetic acid, at a flow rate of 2 ml per minute at the time of sample injection. Ten minutes

- 12 -

following the sample injection, the extract is eluted from the column with a linear gradient of acetonitrile (30% to 65%) in water (70% to 35%) containing 0.05% acetic acid which increases over a 120 minute period. The target compound elutes as a single UV-absorption peak approximately 56 minutes following injection (Figure 21). As discussed in the examples which follow, this isolated and purified compound has a molecular weight of either 348 or 353 amu.

Another embodiment of the invention is directed to another isolated and purified compound derived from an extract of *Zea mays*. As with the previous compound, a *Zea mays* extract is prepared according to the methods of the present invention. The isolated and purified compound is then purified from the extract using a semipreparative C<sub>18</sub> reversed phase HPLC column equilibrated in 70% water and 30% acetonitrile containing 0.05% acetic acid, at a flow rate of 2 ml per minute at the time of sample injection. Ten minutes following the sample injection, the extract is eluted from the column with a linear gradient of acetonitrile (30% to 65%) in water (70% to 35%) containing 0.05% acetic acid which increases over a 120 minute period. The target compound elutes as a single UV-absorption peak approximately 60 minutes following injection (Figure 21). As discussed in the examples which follow, this isolated and purified compound has a molecular weight of either 348 or 353 amu.

Although chromatographic behavior varies slightly from system to system, one reasonably skilled in the art can isolate the above two compounds from a C<sub>18</sub> HPLC column, by eluting the two compounds in a fashion similar to that described above.

Another embodiment of the invention is directed to an isolated and purified compound derived from an extract of *Zea mays* having a molecular weight of 348 amu. In a preferred embodiment, the extract is derived from *Zea mays* corn cob using a polar solvent according to the process of the present invention, and as described in more detail in the examples which follow.

Another embodiment of the invention is directed to an isolated and purified compound derived from an extract of *Zea mays* having a molecular weight of 353 amu. In a preferred embodiment, the extract is derived from *Zea mays* corn cob

- 13 -

using a polar solvent according to the process of the present invention, and as described in more detail in the examples which follow.

The following examples illustrate embodiments of the invention, but should not be viewed as limiting the scope of the invention.

## 5 Examples

### Example 1 Effect of a Classical Phytoestrogenic Type Substance on Uterine Wet Weight.

The uterine wet weight (white bar) and DNA content (stippled bar) of ovariectomized rats housed on wood chips or in wire cages were determined and are  
10 presented in Figure 1. There were ten animals in each group and data are plotted as the mean  $\pm$  SEM for 10 experimental observations. By 14 days following ovariectomy castration, uterine wet weight (66 mg) and DNA content (460 mg) regressed to basal levels (data not shown) for the animals housed on wire. However, both uterine wet weight (95 mg) and DNA content (670  $\mu$ g) were increased by nearly 45% in animals  
15 housed on the wood chip bedding. These data indicated that the wood chip bedding contained a substance with "estrogenic" activity capable of stimulating uterine hypertrophy and hyperplasia beyond that observed in the ovariectomized rats housed on wire.

### 20 Example 2 Effects of Housing Conditions on Uterine Wet Weight and DNA Content

To demonstrate the effects of "phytoestrogen-like substances" on the rat uterus, adult 45 day old Sprague Dawley Rats (Holtzman, Madison, Wisconsin) were either left with ovaries intact (INTACT) or ovariectomized (OVEX) and the animals housed for 12 days in wire cages (INTACT WIRE and OVEX WIRE) or on corn cob  
25 bedding (INTACT CORN and OVEX CORN) suspected of containing phytoestrogens. There were six animals in each of the groups. Following a 12 day period, uterine wet weight (Figure 2A) and DNA content (Figure 2B) were determined by standard procedures (B.M. Markaverich et al., Endocrinology. 114, 814-820, 1984; B.M. Markaverich et al. Environ. Health Perspect. 103: 574-581, 1995; M. Marcelli et al.  
30 Endocrinology 136, 1040-1048, 1995). The data in Figure 2A demonstrate that

- 14 -

regardless of whether the animals were housed on wire or corn cob bedding, ablation of endogenous estrogen by ovariectomy resulted in a dramatic decrease in uterine wet weight (compare INTACT WIRE to OVEX WIRE and INTACT CORN to OVEX CORN). These data suggested the ground corn cob bedding (GCCB) did not contain  
5 significant quantities of phytoestrogenic-like substances, as was the case for the wood chip bedding of Example 1 above.

Analysis of uterine DNA content from the animals in this study also supported this contention and are presented in Figure 2B. As seen in Figure 2B, ovariectomy resulted in a significant decrease in uterine DNA content in animals housed  
10 on wire and a more dramatic decrease was observed in the uteri from animals housed on GCCB. These finding suggested that these animals were absorbing or ingesting a component(s) from corn cob bedding with anti-estrogenic, progestational or perhaps glucocorticoid-like activity. When administered acutely, the weaker estrogens such as estriol and estradiol-17 $\alpha$  can be antagonists of more potent estrogens such as estradiol.  
15 However, sustained exposure (as is the case for the present studies) to these impeded estrogens causes maximum uterotrophic response (J.H. Clark et al., J. Steroid Biochem. 16: 323-328, 1982; J.H. Clark et al., J. Steroid Biochem. 1005-1013., 1984; B.M. Markaverich et al., Endocrinology. 105: 1458-1462. 1979) and can cause mammary cancer (R.L. Noble et al., Cancer Research 35, 766-780, 1975). Therefore, if the GCCB  
20 were releasing weak or impeded estrogens, sustained exposure to this bedding material should have caused hyperestrogenization of the uterus (as in Example 1) and this was not observed. These studies indicate that the "active" substance in corn cob bedding which is affecting mammalian reproduction is not a classical estrogen or phytoestrogen.

Example 3     Effects of GCCB on the Estrous Cycle of Adult Female Harlan Sprague  
25                     Dawley Rats.

Adult Sprague Dawley rats (5 animals per group) were housed in either wire cages or on GCCB. The estrous cycles were monitored by recording daily vaginal smears, presented in Figure 3, for periods of up to either two (black bar) or five (white bar) weeks. 85-90% of the animals housed on wire displayed normal 4-5 day estrous  
30 cycles. Significantly, the numbers of animals that displayed normal 4-5 day cycles was



- 15 -

decreased by 50% in the case of GCCB exposure for two weeks and by 80% in the case of GCCB exposure for five weeks compared to that of the control groups housed on wire for either two or five week periods. Careful inspection of the vaginal smears from the animals housed on GCCB indicated that the majority of these animals were in persistent metestrus which would impede breeding efficiency.

Example 4     Reversibility of the GCCB-Induced Effects on the Estrous Cycle of Adult Female Harlan Sprague Dawley Rats.

The data presented in Figure 4 reveal that the effects of GCCB on the estrous cycles of adult female rats is reversible. For this study, vaginal smears were taken daily for a group of ten adult Harlan Sprague Dawley rats that were housed first in wire cages for 14 days (WIRE), transferred to GCCB for 14 days (CORN) and then returned to wire cages for 14 more days (CORN TO WIRE). The numbers of animals displaying normal 4-5 day estrous cycles reveal that the effects of corn cob bedding on the estrous cycles of these animals was reversible. Again, careful inspection of the vaginal smears from the animals housed on GCCB indicated that the majority of these animals were in persistent metestrus which would impede breeding efficiency.

Example 5     Effects of GCCB on Estrogen Receptor and Type II binding sites.

The effects of GCCB on estrogen receptor binding sites and Type II receptor binding sites were assessed. The results of the analysis are shown in Figures 6A and 6B. Briefly, cytosolic fractions (Figure 6A, estrogen receptors) or nuclear fractions (Figure 6B, Type II sites) were prepared by standard procedures. GCCB was extracted with methanol, dried under nitrogen and resolubilized (1:1000) in 10% ethanol in 10 mM Tris, 1 mM EDTA buffer (crude extract). Alternatively, the 80% methanol in water eluate of an aliquot of GCCB extract was purified via chromatography on a Spice C<sub>18</sub> cartridge (Pur. Extract, closed circles) dried and also rediluted (1:1000) in 10 mM Tris, 1 mM EDTA 10% ethanol buffer. Aliquots of the rediluted extract (1-50 µl), ethanol-TE buffer vehicle only (Control) or the indicated concentration of the estradiol competitors diethylstilbestrol (DES, closed squares), daidzein (open triangles), genistein (closed triangles), or coumestrol (open squares) were added to cytosolic (ER) or nuclear preparations (Type II sites) containing [<sup>3</sup>H]

- 16 -

estradiol (10nM for estrogen receptor; 30 nM for Type II) and binding to estrogen receptor or Type II sites was determined by the method of Markaverich et al. (Environ. Health Perspect. 103: 574-581, 1995; Endocrinology 109: 62-69, 1981).

The results demonstrate that unlike classical phytoestrogens such as daidzein, genistein, and coumestrol, which competitively inhibit [3H] estradiol binding to the estrogen receptor (Figure 6A), neither the crude nor purified corn bedding extract competed for ER binding in rat uterine cytosol over the limited solubility range used. The results suggest that these GCCB extracts do not contain significant quantities of a compound which interacts with the hormone binding domain of the estrogen receptor.

10 Example 6 Effects of GCCB on Female and Male Sexual Behavior.

Exposure to GCCB also had significant effects on male and female mating behavior, shown in Figures 5A and 5B. Briefly, ovariectomized female rats were primed subcutaneously with 2 µg of estradiol benzoate in sesame oil. Forty-eight hours later they were administered 100 µg of progesterone subcutaneously. Four hours after progesterone administration, their sexual behavior was analyzed in the presence of male proven breeder rats. Sexual behavior was scored and quantitated as lordosis response. The females were on Sani-chip bedding (control) or on GCCB (corn-cob). The results are presented in Figure 5A.

Adult male proven breeders were placed in individual cages on Sani-chip bedding (control, open circles) or GCCB (corn-cob, closed circles) and their sexual performance evaluated in the presence of sexually receptive ovariectomized, hormone-treated female rats. Male sexual behavior was evaluated by examining number of mounts, intromissions, ejaculations, ejaculation latencies (in seconds) and grooming frequencies in a 30-minute test period with each sexually receptive female. The results are listed in Figure 5B.

When female mating behavior of rats housed on GCCB was monitored for over 11 weeks, only 10-15% of the animals displayed lordosis as compared to the Sani-chip controls (Figure 5A). Similarly, housing of male animals on GCCB significantly blocked male sexual behavior (Figure 5B). Mounting, intromission and ejaculation efficiencies were decreased by nearly 99% in GCCB housed males relative

- 17 -

to Sani-chip controls. Ejaculation latency and grooming frequency were also decreased by 75-80% in animals housed on GCCB (Figure 5B). These observations demonstrate that GCCB contains an activity which is capable of affecting male and female reproductive function.

5    Example 7    Extraction of Corn Cob and GCCB.

GCCB, fresh corn, kernels, and cobs were extracted separately in 100% methanol at 70°C for 2 hours. The extracts were dried under vacuum at 70°C and redissolved in dimethyl sulfoxide (DMSO). Aliquots of the redissolved extract (2, 5 or 10 µl) were added to cultured MCF-7 human breast cancer cells.

10   Example 8    Isolation of a Mitogenic Activity from GCCB Extract.

An aliquot of the GCCB crude extract was fractionated on a Spice C<sub>18</sub> Cartridge (a reverse phase column using aliphatic groups attached to a matrix to form a hydrophobic adsorbent. Rainin Instruments Company). Briefly, a GCCB extract in 20% methanol in water was loaded onto a spice cartridge and the pass through (20%  
15    methanol) fraction was collected. The column was sequentially eluted with a solution of water comprising 40%, 60%, and 80% methanol in water. The final elution was with 100% methanol. Each fraction, including the pass through fraction, was collected and dried at 50°C under nitrogen to remove methanol. The eluate was re-dissolved in 100% ethanol. Aliquots of each fraction were added to cultures of MCF-7 (ER+) or MDA-  
20    231 (ER-) human breast cancer cells grown in phenol red free medium. Control wells received only the ethanol vehicle. Twenty-four hours after plating the cells, the media was changed and the cells were treated with various doses of GCCB extract (2-10 ml) alone, or in combination with 10 nM estradiol. Compounds were added to the media in ethanol and ethanol concentration was maintained at 0.2% in all wells including  
25    controls. Six days following treatment, cell number was determined by hemocytometer counts and results are expressed as cell number in treated wells relative to ethanol controls (100%) as described (B.M. Markaverich et al., J. Steroid Biochem. 29:71-78, 1988; B.M. Markaverich et al., Cancer Res. 50: 1470-1478, 1990; B.M. Markaverich et al., JBC 263: 7203-7210, 1988; B.M. Markaverich et al., Internat. J. Oncol. 4:1291-  
30    1300, 1994).

- 18 -

The results of this study are shown in Figure 7. The crude extract stimulated the proliferation of MCF-7 cells (black bars), but not MDA-231 cells (white bars). All of the mitogenic activity was recovered in the 80% methanol in water eluate from the Spice C<sub>18</sub> cartridge. The observation that the corn cob bedding extract failed to stimulate the proliferation of ER-negative MDA-231 human breast cancer cells is of interest. At higher concentrations, in the presence or absence of estradiol, corn cob extract inhibited growth of MDA-231 cells. The differential response of MCF-7 and MDA-231 cells to corn cob extract suggests that the ER-regulated pathways are at least peripherally involved in the mechanism through which the active component in the corn cob extract stimulates MCF-7 cell proliferation.

In a further study, the effects of fractionated GCCB extract on the proliferation of the MCF-7 and MDA-231 human breast cancer cells were analyzed. Briefly, a crude GCCB extract was prepared as described above and diluted 1:1000 in TE methanol and fractionated on a Spice C<sub>18</sub> cartridge. Briefly, fractionation was accomplished by loading the extract (2 ml) onto the column and collecting the pass through (20% methanol in water) fraction. The column was then successively eluted with 2 ml volumes of 40%, 60%, 80% methanol in water solutions. Final elution was with 100% methanol. Each of these fractions were collected for assay. Aliquots of the crude extract (Whole Ext) and the 20%, 40%, 60%, 80% and 100% methanol cartridge eluates were taken to dryness at 50°C under nitrogen, redissolved in 500 µl of ethanol and 20 µl aliquots were added to the cells 24 hours following plating (Day 0). Six days following treatment, cell number (% Control) was determined. Data represent the mean SEM and were analyzed statistically via ANOVA and Tukeys test on the means using Instat. As can be seen in Figure 8, the response of MCF-7 cells (black bars) to the purified material (80% methanol fraction) was equivalent to that obtained with the crude extract or estradiol.

Experimental studies were performed to determine if GCCB extract has an effect on [<sup>3</sup>H]estradiol binding to estrogen receptor or nuclear Type H sites in rat uterine preparations. Cytosolic and nuclear fractions were prepared by standard procedures (B.M. Markaverich et al., Environ. Health Perspect. 103: 574-581, 1995; M.

- 19 -

Marcelli et al. Endocrinology 136, 1040-1048, 1995). GCCB was extracted with methanol, and the crude extract dried under nitrogen and rediluted (1:1000) in 100% ethanol in TE buffer (crude extract). Alternatively, the 80% methanol in water eluate of an aliquot of GCCB extract was purified via chromatography on a Spice C<sub>18</sub> Cartridge (Pur. Extract), dried and also rediluted (1:1000) TE-10% ethanol buffer. Aliquots of the rediluted extract (1-50 µl), ethanol-TE buffer vehicle (control) or the indicated concentrations of diethylstilbestrol (DES), daidzein, genistein, or coumestrol were added to the cytosolic (estrogen receptor) or nuclear preparations (Type II sites) containing [<sup>3</sup>H]estradiol (10 nM for estrogen receptor, 30 nM for Type II) and binding to estrogen receptor or Type II sites was determined by well documented procedures (Markaverich, B.M. et al. Environ. Health Perspect. 103: 574-581, 1995; B.M. Markaverich et. al. Endocrinology. 109: 62-69, 1981). As seen in Figure 6, GCCB extract competed for [<sup>3</sup>H]estradiol binding to nuclear Type II sites, but not to estrogen receptors.

Further characterization of the 80% methanol eluate from the Spice cartridge by HPLC revealed that the GCCB extract eluted from a Waters microbondapack C<sub>18</sub> column as a single major peak of UV absorbing material. Fractions were collected as indicated in the chromatogram and aliquots were added to MCF-7 human breast cancer cell cultures. The mitogenic activity in GCCB extract eluted from a C<sub>18</sub> Sephadex column as a single, major component. The data in Figure 10 demonstrate that a single peak of mitogenic activity eluted from the column together with a peak of UV absorbing material. The symmetry of the HPLC peak suggests that this material has been purified to near homogeneity and that the compound of Figure 9 shown to interact with Type II sites is likely to be responsible for stimulating MCF-7 human breast cancer cell proliferation. This stimulation was not observed in MDA-231 cells and, in fact, the preparations partially inhibited MDA-231 cell proliferation. This is similar to the results achieved with partially purified preparations and the HPLC purified material, shown in Figures 7 and 8.

Example 9     Effects of Cob Bedding Extract on ER-Mediated Gene Transcription.

- 20 -

The effect of purified ground corn cob bedding extract on estrogen mediated gene transcription in mammalian cells was determined using HeLa cells lacking endogenous estrogen receptors (ER). The ER-negative HeLa cells were co-transfected with the construct psvmt:wE, which encodes for exogenous estrogen receptor, the construct pERE-E1b-LUC, which contains an estrogen responsive element (ERE) regulating the expression of the luciferase reporter gene and the construct pCMV $\beta$ gal, which encodes a housekeeping gene to assess basal levels of gene expression in these cells. Expression of pCMV $\beta$ gal was used for normalization of the response of the pERE-E1b-LUC reporter construct to estradiol and the ground corn cob bedding extract.

As shown in Figure 11, treatment of HeLa cells with either 1 nM estradiol (E<sub>2</sub>) or 10 nM of the phytoestrogen genistein (GEN) resulted in a stimulation of ER-mediated luciferase activity greater than two fold of that of the control (EtOH). Treatment of the transfected cells with either a crude ground corn cob bedding extract (CRUDE X) or purified ground corn cob bedding fractions eluting from Spice C<sub>18</sub> Cartridges (20%, 40%, 60%, 80% or 100% methanol eluates) failed to significantly stimulate the activity of the reporter transgene relative to the control. These data indicate that it is unlikely that a component in the ground corn cob bedding extract affects transcription via direct binding interactions with the estrogen receptor or through an estrogen-responsive signal transduction pathway.

Example 10 Effects of Anti-estrogen, Estrogen or Corn Cob Bedding Extract Stimulation of MCF-7 Cells.

Experimental studies were performed to assess the ability of a pure anti-estrogen (ICI-182,780) to effect the ability of the ground corn cob bedding extract to stimulate MCF-7 human breast cancer cell proliferation. It is known that antiestrogen ICI-182,780 blocks cellular proliferation at the level of the estrogen receptor. If the ability of the ground corn cob bedding extract to stimulate MCF-7 cell proliferation (see Figures 7, 8 and 10) is mediated via direct interaction with the estrogen receptor or through an estrogen-responsive signal transduction pathway, then one would predict

- 21 -

that ICI-182,780 would block this corn cob extract-dependent proliferative response of MCF-7 cells.

Exponentially growing cultures of MCF-7 cells were treated with 1 nM estradiol ( $E_2$ ), 10 nM ICI-182,780 (ICI) or 10  $\mu$ l of the highly purified ground corn cob bedding extract (80% eluate from Spice  $C_{18}$  Cartridge) alone or in combination. Cell number was determined 6 days following treatment. As shown in Figure 12,  $E_2$  treatment resulted in a 2-3 fold stimulation of MCF-7 cell proliferation relative to control (CONT). Although ICI-182,780 (ICI) alone failed to inhibit the proliferation of these cells in the absence of estrogen, antagonistic activity of ICI-182,780 was observed when it was added to cells stimulated with  $E_2$  (compare  $E_2$  versus ICI versus  $E_2$  + ICI treatment groups). The ICI antiestrogen blocked greater than 50% of the estradiol-induced stimulation of MCF-7 breast cancer cell proliferation.

In addition, the purified ground corn cob bedding extract (EXT) alone stimulated MCF-7 cell proliferation (Figure 12) and this response was not blocked by the antiestrogen (compare EXT group to ICI + EXT group). These data suggest that the active mitogen in the ground corn cob bedding extract is not a typical phytoestrogen, mycotoxin or xenobiotic which controls cellular proliferation via direct interaction with the estrogen receptor. Thus, the mitogenic agent in ground corn cob bedding may control cellular growth and proliferation by a previously unknown mechanism.

#### 20 Example 11 Effects of Highly Purified Corn Cob Bedding Extract on the Estrous Cycle.

Experimental studies were performed to determine if the material isolated from the highly purified corn cob bedding extract is responsible for the disruption of normal estrous cycles in animals. Ground corn cob was extracted as described in Example 8. The extract was purified by chromatography on Spice  $C_{18}$  cartridges. The 80% methanol eluate was examined to determine whether it had biological activity in animal studies.

Thirty adult female rats housed in wire cages were divided into three treatment groups (controls, low dose extract and high dose extract groups) of ten rats each. Food and water were provided *ad libitum* and the estrous cycles of these animals

- 22 -

were followed by monitoring daily vaginal smears for 6 weeks. The low dose group was fed 50  $\mu$ l of the 80% methanol eluate per 400 ml drinking water. The high dose group was fed 100  $\mu$ l of the 80% methanol eluate per 400 ml of drinking water. As shown in Figure 13, the results of the study demonstrate that animals in all three groups have normal 4-5 day estrous cycles for the initial 6 weeks of the study.

At the end of the sixth week, the animals in these three groups were given either a low dose (400  $\mu$ l corn cob extract per 400 ml water) or high dose (2000  $\mu$ l extract per 400 ml water) of ground corn cob bedding extract in the drinking water or water only (controls) and fluid consumption was monitored twice weekly throughout the study. The methanol solvent in the corn cob extract was replaced with ethanol. Aliquots of the 80% methanol extract were dried under nitrogen, redissolved in equal volumes of 100% ethanol and added to the drinking water in this vehicle. To control for any ethanol-dependent effects, a volume of ethanol equivalent to the high dose (2000  $\mu$ l per 400 ml of drinking water) was also added to the drinking water as control. No effects on body weights, fluid consumption or estrous cycle duration were noted as a result of treatment with the ethanol vehicle control.

Daily vaginal smears were collected from animals in all three groups for an additional 3 weeks. The data clearly show that 70% of the animals receiving the low dose ground corn cob bedding extract displayed abnormal estrous cycles following 1 week of treatment and 100% of these animals were acyclic by 2 or 3 weeks following treatment. A more pronounced effect was observed with the high dose of ground corn cob bedding extract where 100% of the animals were acyclic during all three weeks of the treatment period. These data demonstrate that the highly purified ground corn cob bedding extract contains a substance(s) which interrupts normal cyclicity in female rats.

#### Example 12 High Performance Liquid Chromatography of GCCB Extract.

An aliquot of the 80% methanol fraction of GCCB extract was injected onto a C<sub>18</sub> (Waters micro-bondapak) reversed phase column eluted with 70% methanol in water at a flow rate of one ml per minute. One ml fractions were collected (one each minute) and taken to dryness under nitrogen at 50°C, resuspended in 10  $\mu$ l of 100% ethanol. The fractions were added to exponentially growing MCF-7 cells and also the



- 23 -

O.D.<sub>280</sub> absorbance was determined for each fraction. Cell number was determined 6 days following treatment and data are presented in Figure 10 as the mean  $\pm$  the SEM for twelve determinations. The cell number determined for each fraction (% Control) is shown plotted over the O.D.<sub>280</sub> absorbance tracing representing the chromatogram. The data reveal that a single peak of MCF-7 stimulating activity (closed circles) eluted from the column together with a single peak of UV absorbing material. Thus, the UV absorption activity co-elutes with cell proliferation stimulating activity.

Example 13 Effects of Purified GCCB Extract on [<sup>3</sup>H]estradiol Binding to Estrogen Receptor or Nuclear Type II Sites.

Rat uterine cytosolic fractions (estrogen receptor) and nuclear fractions (Type II sites) were prepared as follows. The 80% methanol in water eluate of an aliquot of GCCB extract purified via chromatography on a Spice C<sub>18</sub> Cartridge was dried and rediluted (1:1000) TE-10% ethanol buffer. Aliquots of the rediluted extract (1-50  $\mu$ l) in ethanol-TE buffer vehicle (control) were added to the cytosolic and nuclear fractions containing [<sup>3</sup>H]estradiol (10 nM for estrogen receptor, 30 nM for Type II sites) and binding to estrogen receptor (ER) or Type II sites (TYPE II) was determined by well documented procedures (B.M. Markaverich et al. Environ. Health Perspect. 103: 574-581, 1995; B.M. Markaverich et al. Endocrinology 109: 62-69, 1981). The results in Figure 9 show that highly purified GCCB extract preparations competed for [<sup>3</sup>H]estradiol binding to nuclear Type II sites, but did not compete for binding to the estrogen receptor.

Example 14 Effects of diluted Extract on Estrogen Receptor Negative Cells.

Experimental studies were performed to determine if the corn cob extract at a more dilute concentration comprises mitogenic activity. It is possible that MDA-231 cells are not stimulated by the corn cob bedding because they may be more sensitive to the mitogenic agent than are MCF-7 cells. Shown in Figure 14, 5 ml of various dilutions (1, 2, 5, 10, 50 and 100 fold dilutions) of corn cob extract were added to exponentially growing MDA-231 cells. Cell number was determined by hemocytometer counting 11 days following treatment. The bar denoted 0 is data from MDA-231 cells treated with 5 ml of the ethanol vehicle as a control. While

- 24 -

concentrated preparations of the corn cob extract inhibited MDA-231 cell proliferation, cell stimulation became apparent as the corn cob extract was diluted (compare 10, 50 and 100 fold dilutions with the 1, 2 and 5 fold dilutions).

In a separate experiment, exponentially growing MDA-231 cells were  
5 treated with 2  $\mu$ l (grey bars), 5  $\mu$ l (black bars), 10  $\mu$ l (striped bars) of the indicated dilution (1, 2, 5, 10, 50 and 100 fold) of a corn cob bedding extract in ethanol or ethanol vehicle only as a control (bar labeled 0). Cell number was determined by hemocytometer counting 11 days following treatment. As seen in Figure 15, 2  $\mu$ l and 5  $\mu$ l of a 10 fold, 50 fold, or 100 fold dilution of corn cob extract stimulated the  
10 proliferation of MDA-231 human breast cancer cells. These data clearly indicate that MDA-231 cell growth inhibition is observed with higher concentrations of these extracts and mitogenic activity is observed with lower concentrations of the extract.

Example 15 Effects of Methotrexate and GCCB Extracts on Breast Cancer Cells.

Exponentially growing MCF-7 human breast cancer cells were treated  
15 with Methotrexate alone (open and closed circles), Methotrexate and estradiol (E2, open squares), or a combination of Methotrexate and purified corn cob bedding extract (Ext, closed squares). Cell number was determined 11 days following treatment. Vehicle controls (DMSO, open circles and ethanol, closed circles) were added to the Methotrexate alone in order to control for any vehicle-dependent effects in the  
20 experiment. The experimental data are shown in Figure 16. The combination of Methotrexate and corn cob extract suppressed the proliferation of MCF-7 cells more effectively than Methotrexate and estradiol.

Example 16 Effects of Fresh Corn Extract on Cell Growth.

Fresh corn on the cob was purchased at a local supermarket and the  
25 kernels were separated from the cob. Both the kernel and the cob were separately extracted by the method of Example 7. Briefly, extraction was performed in 100% methanol for 2 hours at 70°C. The extract was dried under vacuum and redissolved in DMSO and aliquots of about 2  $\mu$ l, 5  $\mu$ l and 10  $\mu$ l were added to cultured MCF-7 cells. After 2 days, the cells were counted. The data is plotted in Figure 17. Low doses of  
30 either the fresh kernel (Kern) or fresh cob (Cob) extract stimulated MCF-7 human

- 25 -

breast cancer cell proliferation relative to controls whereas higher doses of these preparations inhibited cell proliferation. A classical bell-shaped dose response curve was obtained for both the corn cob and corn kernel preparations. This demonstrates that extract from fresh corn on the cob contains a mitogenic activity similar to that described  
5 for extract from ground corn cob bedding.

Example 17 Further Purification of GCCB Extract by High Performance Liquid Chromatography.

An aliquot of the 80% methanol fraction of GCCB extract was injected onto a semi-preparative Altex Ultrasphere-ODS reversed phase column HPLC column  
10 (10 mm ID x 25 cm length) equilibrated in 70% Solvent A (HPLC grade water containing 0.05% acetic acid) and 30% Solvent B (HPLC grade acetonitrile containing 0.05% acetic acid). The solvent flow rate through the column was 2 ml per minute. Ten minutes following injection of the GCCB preparation, a linear gradient increasing to 65% Solvent B over a 120 minute interval was initiated and the run was continued  
15 for 140 minutes. Four minute fractions of the column eluate were collected from injection (time 0) through 140 minutes. Aliquots of the fractions were dried, redissolved in ethanol and added to exponentially growing MCF-7 human breast cancer cells (large filled circles) and cell number was determined 6 days following treatment. The data shown in Figures 18A and 18B suggested that the MCF-7 stimulating activity  
20 eluted from the column coincident with two very small peaks of UV-absorbing material. In Figure 18A, the mitogenic activity of Fraction 14 (arrow labeled Fr 14) was not resolved into two components because of the large (four minute interval) fraction size. To confirm multiple components, an aliquot of Fraction 14 was re-injected onto a C<sub>18</sub> Ultrasphere-ODS column and eluted as described above to check purity. Re-  
25 chromatography of Fraction 14, indicated by the solid line in Figure 18B, revealed that Fraction 14 consisted of two very small UV-absorbing peaks (arrow labeled Fr 14 in Figure 18B), which co-eluted with the mitogenic activity.

- 26 -

Example 18    Molecular Weight Determination of the Putative Mitogenic Agents in Corn Cob Bedding Extracts by Mass Spectrometry.

An aliquot of the HPLC purified Fraction 14 was diluted in an aqueous mobile phase consisting of 50% acetonitrile containing 4 mM ammonium acetate. This was loop injected at 5  $\mu$ L per minute onto an API 3000 triple quadrupole mass spectrometer (PE Siex Concord, Ontario, Canada) for mass spectrometric (MS) and tandem mass spectrometric (MS/MS) analysis. Samples were ionized under atmospheric pressure ionization conditions with a skimmer voltage of 70V and the instrument was tuned to detect positive ions. The first quadrupole was scanned to detect ions for MS analysis or held static to transmit selected ions for MS/MS analyses. Collisionally activated decomposition of selected ions was achieved in the rf-only quadrupole collision cell with nitrogen as the collision gas and collision energies between 30 and 60 eV. The third quadrupole was scanned to detect positive ions. The results of this analysis confirmed HPLC studies demonstrating that Fraction 14 contained two substances. The molecular weights of these substances were determined to be 348 amu (Figure 19) and 353 amu (Figure 20), respectively. Since these were the only components detected in the preparations, both substances have a high probability of representing the active agents in the GCCB preparations. More sophisticated analysis by gas chromatography-mass spectrometry (GC-MS), infrared spectrometry (IRS) and nuclear resonance spectrometry (NMRS) may be used to determine the chemical structure(s) of the 348 and 353 amu components in the GCCB samples and establish these compounds as the mitogenic agents in cultured breast cancer cells and as the biological activity in behavioral studies with male and female animals studies described above.

Example 19    Confirmation of Two Mitogenic Agents in GCCB Extracts by HPLC.

A sample of GCCB was prepared and chromatographed on HPLC under conditions identical to those described in Example 17 with the exception that fractions were collected at one minute intervals instead of four minute intervals. The smaller fraction size increased the resolution of the components four fold as opposed to pooling fractionated material over four minute intervals as was done in Example 17. Aliquots

- 27 -

of each of the fractions were assayed for mitogenic activity in MCF-7 human breast cancer cells as was described for Examples 7, 8 and 10. The data, shown in Figure 21, clearly demonstrate that two peaks of mitogenic activity eluted from the column coincident with corresponding peaks of UV-absorbing material. These two agents  
5 likely represent the 348 and 353 molecular weight components observed by HPLC-MS analysis, shown in Figures 19 and 20, respectively, and are strong candidates for the active components in the GCCB extract preparations.

Other embodiments and uses of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention  
10 disclosed herein. All U.S. and foreign patents and patent applications, including the provisional patent application, serial number 60/087,680, filed June 2, 1998, and all other documents referenced herein, for whatever reason, are specifically incorporated by reference. The specification and examples should be considered exemplary only with the true scope and spirit of the invention indicated by the following claims.

15

- 28 -

We Claim:

1. A process for producing a therapeutic substance comprising:  
extracting a *Zea mays* plant product in a first solvent to produce a solvent  
extract;  
5 drying said solvent extract to produce an extracted solid;  
solubilizing said extracted solid in a second solvent;  
purifying said solubilized extract in a chromatographic process;  
collecting an active fraction from said chromatographic process.
2. The process of claim 1 wherein said *Zea mays* plant product is selected from the  
10 group consisting of corn kernel, corn cob, animal bedding material made from  
corn cob, and mixtures thereof.
3. The process of claim 2 wherein said animal bedding material is ground corn cob  
bedding.
4. The process of claim 1 wherein said first solvent is selected from the group  
15 consisting of alcohols, ethers, ketones, water and mixtures thereof.
5. The process of claim 4 wherein said alcohol is methanol.
6. The process of claim 1 wherein said first solvent and said second solvent are the  
same.
7. The process of claim 1 wherein said extracting step is from 1 hour to 1 month.
- 20 8. The process of claim 1 wherein said extraction step is carried out at a  
temperature between 0° C. and 100° C.
9. The process of claim 1 wherein said chromatographic process is a procedure  
selected from the group consisting of column chromatography, batch  
chromatography, precipitation, specific adsorbent chromatography, gel  
25 filtration, and combinations thereof.
10. The process of claim 9 wherein said column chromatography is a reverse phase  
chromatography column comprising a C<sub>18</sub> solid phase and a methanol/water  
liquid phase.
11. The process of claim 1 wherein said active fraction is a fraction of about 80%  
30 methanol.

- 29 -

12. The therapeutic substance produced by the process of claim 1.
13. The therapeutic substance of claim 12, in a form selected from the group consisting of a salt, a compound, a complex and a combination thereof.
14. A pharmaceutical preparation comprising the therapeutic substance produced  
5 by the process according to claim 1 and a pharmaceutically acceptable carrier.
15. The pharmaceutical preparation of claim 14 prepared in the form of an aqueous solution.
16. The pharmaceutical preparation of claim 15 wherein said aqueous solution is for oral use or parenteral use.
- 10 17. The pharmaceutical preparation of claim 14, prepared in the form of capsules for oral use.
18. The pharmaceutical preparation of claim 14, prepared in the form selected from the group consisting of unguents, ointments, creams, suppositories and ovules.
19. A method for treating a patient with a neoplastic disorder comprising  
15 administering to the patient an antineoplastic composition comprising an effective quantity of the therapeutic substance of claim 1.
20. The method of claim 19 wherein said antineoplastic composition further comprises an antineoplastic compound.
21. The method of claim 20 wherein said antineoplastic compound is methotrexate.
- 20 22. The method of claim 20 wherein said antineoplastic compound is selected from the group consisting of 5-fluorouracil, 6-mercaptopurine, 6-thioguanine, alkylating agents, androgens, antiadrenals, antiandrogens, antiestrogens, antimetabolites, asparaginase, bleomycin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, corticosteroids, cyclophosphamide, cytarabine,  
25 dacarbazine, dactinomycin, daunorubicin, doxorubicin, estrogens, etoposide, fludarabine, fluorouracil, flutamide, hexamethylmelamine, hormones, hydroxyurea, hypothalamic agents, idarubicin, ifosfamide, intrapleural, lomustine, mechlorethamine, melphalan, methotrexate, mithramycin, mitomycin, mitotic inhibitors, mitoxantrone, paclitaxel, procarbazine,

- 30 -

progestins, streptozocin, taxol, thiotepa, vinblastine, vincristine, vinorelbine and combinations and analogs thereof.

23. The method of claim 19 wherein said neoplastic disorder is selected from the group consisting of prostate cancer, breast cancer, lymphoma, leukemia, testicular cancer, gastrointestinal tract cancer, lung cancer, skin cancer, head tumors, neck tumors, bone cancer, liver cancer, and pancreatic cancer.
24. A composition for effecting contraception in a patient, said composition produced by the process of claim 1.
25. The composition of claim 24 wherein the patient is a mammal.
26. The composition of claim 24 wherein the patient is a male or a female.
27. The composition of claim 24 wherein the patient is human.
28. An isolated and purified compound derived from the therapeutic substance of claim 1, said compound having a molecular weight of 348 amu.
29. The compound of claim 28 in which the plant product is corn cob.
30. An isolated and purified compound derived from the therapeutic substance of claim 1, said compound having a molecular weight of 353 amu.
31. The compound of claim 30 in which the plant product is corn cob.
32. An isolated and purified compound derived from the therapeutic substance of claim 1, said compound eluting as a single UV-absorption peak from a C<sub>18</sub> reversed phase HPLC column at a rate of 2 ml per minute, said compound eluting approximately 56 minutes after injection of said therapeutic substance into said column, said compound having a molecular weight of either 348 or 353 amu.
33. An isolated and purified compound derived from the therapeutic substance of claim 1, said compound eluting as a single UV-absorption peak from a C<sub>18</sub> reversed phase HPLC column at a rate of 2 ml per minute, said compound eluting approximately 60 minutes after injection of said therapeutic substance into said column, said compound having a molecular weight of either 348 amu or 353 amu.



- 31 -

34. An isolated and purified compound derived from an extract of *Zea mays* having a molecular weight of 348 amu.
35. The compound of claim 34 in which the extract is derived from *Zea mays* corn cob using a polar solvent.
- 5 36. An isolated and purified compound derived from an extract of *Zea mays* having a molecular weight of 353 amu.
37. The compound of claim 36 in which the extract is derived from *Zea mays* corn cob using a polar solvent.
38. A method of contraception comprising:
- 10 isolating an extract from a *Zea mays* plant product using a polar solvent; and administering an effective amount of the extract to an individual to effect contraception.
39. The method of claim 38 wherein the polar solvent is an alcohol.
40. The method of claim 38 wherein the individual is a female and contraception is effected by inhibiting estrus, inhibiting the estrous cycle, inhibiting ovulation, inhibiting oogenesis, inhibiting sexual desire, inhibiting zygote implantation or combinations thereof.
- 15 41. The method of claim 38 wherein the individual is a male and contraception is effected by inhibiting sexual desire, inhibiting spermatogenesis, inhibiting erectile function, or combinations thereof.
- 20 42. The method of claim 38 wherein the plant product is selected from the group consisting of corn kernel, corn cob, animal bedding material made from corn cob, and mixtures thereof.
43. The method of claim 38 wherein the individual is male or female.
- 25 44. The method of claim 38 wherein the individual is a mammal selected from the group consisting of humans, monkeys, mice, rats, guinea pigs, rabbits, hamsters, horses, cattle, pigs, sheep, goats, dogs, and cats.
45. The method of claim 38 wherein the extract is administered orally, topically, intramuscularly, intravenously, subcutaneously, vaginally, rectally or by
- 30 pulmonary absorption.

- 32 -

46. The method of claim 38 further wherein the step of isolating further comprises the step of purifying the extract by a chromatographic process before administration.
47. The method of claim 46 wherein the chromatographic process is reverse phase chromatography.
48. A method of inhibiting neoplastic activity comprising the steps of:  
isolating an antineoplastic extract from a *Zea mays* plant product using a polar solvent; and  
administering an effective amount of the antineoplastic extract to an individual.
49. The method of claim 48 further comprising the step of administering an antineoplastic compound.
50. The method of claim 49 wherein the antineoplastic compound is methotrexate and the neoplastic activity is breast cancer cell proliferation.
51. The method of claim 48 wherein the extract is administered orally, topically, intramuscularly, intravenously, subcutaneously, vaginally, rectally, or by pulmonary absorption.

1/21




FIG. 1

2/21




FIG. 2A




FIG. 2B

SUBSTITUTE SHEET (RULE 26)

3/21




FIG. 3

4/21




FIG. 4

5/21




FIG. 5A




FIG. 5B

6/21




FIG. 6A




FIG. 6B



7/21




FIG. 7

8/21




FIG. 8

9/21




FIG. 9

10/21




FIG. 10

11/21




FIG. 11

12/21




FIG. 12

13/21




FIG. 13

14/21




FIG. 14



15/21




FIG. 15

16/21




FIG. 16

17/21




FIG. 17




FIG. 18A





FIG. 18A

SUBSTITUTE SHEET (RULE 26)

19/21



SUBSTITUTE SHEET (RULE 26)

20/21




FIG. 20

SUBSTITUTE SHEET (RULE 26)

21/21




FIG. 21

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/11880

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K35/78

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 03142 A (BIOGAL GYOGYSZERGYAR) 5 March 1992 (1992-03-05) page 6, line 4 - page 12, line 29 -----	1-4

☐

Further documents are listed in the continuation of box C.

☒

Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

24 August 1999

Date of mailing of the international search report

31/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Rempp, G



# INTERNATIONAL SEARCH REPORT

international application No.

PCT/US 99/11880

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 19-27, 38-51  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 19-27,38-51 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

Continuation of Box I.1

Claims Nos.: 19-27,38-51

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/11880

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9203142 A	05-03-1992	CA 2067770 A	25-02-1992
		CS 9102618 A	18-03-1992
		EP 0497952 A	12-08-1992
		JP 5503295 T	03-06-1993
		PT 98757 A	31-07-1992

**THIS PAGE BLANK (USPTO)**